

Welcome to STN International! Enter x:x

LOGINID:sssptal600rxa

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
 NEWS 2 Apr 08 "Ask CAS" for self-help around the clock  
 NEWS 3 Apr 09 BEILSTEIN: Reload and Implementation of a New Subject Area  
 NEWS 4 Apr 09 ZDB will be removed from STN  
 NEWS 5 Apr 19 US Patent Applications available in IFICDB, IFIPAT, and IFIUDB  
 NEWS 6 Apr 22 Records from IP.com available in CAPLUS, HCAPLUS, and ZCAPLUS  
 NEWS 7 Apr 22 BIOSIS Gene Names now available in TOXCENTER  
 NEWS 8 Apr 22 Federal Research in Progress (FEDRIP) now available  
 NEWS 9 Jun 03 New e-mail delivery for search results now available  
 NEWS 10 Jun 10 MEDLINE Reload  
 NEWS 11 Jun 10 PCTFULL has been reloaded  
 NEWS 12 Jul 02 FOREGE no longer contains STANDARDS file segment  
 NEWS 13 Jul 22 USAN to be reloaded July 28, 2002;  
 saved answer sets no longer valid  
 NEWS 14 Jul 29 Enhanced polymer searching in REGISTRY  
 NEWS 15 Jul 30 NETFIRST to be removed from STN  
 NEWS 16 Aug 08 CANCERLIT reload  
 NEWS 17 Aug 08 PHARMAMarketLetter(PHARMAML) - new on STN  
 NEWS 18 Aug 08 NTIS has been reloaded and enhanced  
 NEWS 19 Aug 09 JAPIO to be reloaded August 25, 2002  
 NEWS 20 Aug 19 Aquatic Toxicity Information Retrieval (AQUIRE)  
 now available on STN  
 NEWS 21 Aug 19 IFIPAT, IFICDB, and IFIUDB have been reloaded  
 NEWS 22 Aug 19 The MEDLINE file segment of TOXCENTER has been reloaded  
 NEWS EXPRESS February 1 CURRENT WINDOWS VERSION IS V6.0d,  
 CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP),  
 AND CURRENT DISCOVER FILE IS DATED 05 FEBRUARY 2002  
 NEWS HOURS STN Operating Hours Plus Help Desk Availability  
 NEWS INTER General Internet Information  
 NEWS LOGIN Welcome Banner and News Items  
 NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
 NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 13:02:52 ON 21 AUG 2002

=> fil reg

COST IN U.S. DOLLARS

SINCE FILE  
ENTRY

TOTAL  
SESSION  
0.21

FULL ESTIMATED COST

0.21

FILE 'REGISTRY' ENTERED AT 13:03:00 ON 21 AUG 2002  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 American Chemical Society (ACS)

STRUCTURE FILE UPDATES: 20 AUG 2002 HIGHEST RN 444542-98-7  
DICTIONARY FILE UPDATES: 20 AUG 2002 HIGHEST RN 444542-98-7

TSCA INFORMATION NOW CURRENT THROUGH MAY 20, 2002

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Calculated physical property data is now available. See HELP PROPERTIES  
for more information. See STNote 27, Searching Properties in the CAS  
Registry File, for complete details:  
<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

=>


Uploading 10058456.str

L1 STRUCTURE UPLOADED

=> d

L1 HAS NO ANSWERS

L1 STR



G1 H, X, O, C

G2 C, H, X

Examiner Anderson 703-605-1157

Structure attributes must be viewed using STN Express query preparation.

=> s l1 full

FULL SEARCH INITIATED 13:03:26 FILE 'REGISTRY'  
FULL SCREEN SEARCH COMPLETED - 20061 TO ITERATE


100.0% PROCESSED 20061 ITERATIONS  
SEARCH TIME: 00.00.03

10 ANSWERS

L2 10 SEA SSS FUL L1

=> d scan


L2 10 ANSWERS REGISTRY COPYRIGHT 2002 ACS  
 IN 1H-Imidazol-2-amine, N-[2-bromo-6-chloro-4-(1-methylethyl)phenyl]-4,5-  
 dihydro-, monohydrobromide (9CI)  
 MF C12 H15 Br Cl N3 . Br H  
 CI COM



• HBr

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1


L2 10 ANSWERS REGISTRY COPYRIGHT 2002 ACS  
 IN 1H-Imidazol-2-amine, N-[2-bromo-6-chloro-4-(1-methylethyl)phenyl]-,  
 monohydrochloride (9CI)  
 MF C12 H17 N3 O . Cl H  
 CI COM



• HCl

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1


L2 10 ANSWERS REGISTRY COPYRIGHT 2002 ACS  
 IN 2-Imidazoline, 2-cumidino- (7CI, 8CI)  
 MF C12 H17 N3  
 CI COM



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L2 10 ANSWERS REGISTRY COPYRIGHT 2002 ACS  
 IN 1H-Imidazol-2-amine, N-[2-bromo-6-chloro-4-(1-methylethyl)phenyl]-4,5-  
 dihydro- (9CI)  
 MF C12 H15 Br Cl N3  
 CI COM




\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L2 10 ANSWERS REGISTRY COPYRIGHT 2002 ACS  
IN Butanedioic acid, compd. with N-(4,5-dihydro-1H-imidazol-2-yl)-7-(1-methylethyl)-1H-indol-5-amine (1:1) (9CI)  
MF C14 H18 N4 . C4 H6 O4

CM 1




CM 2



HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1


L2 10 ANSWERS REGISTRY COPYRIGHT 2002 ACS  
IN 2-Imidazoline, 2-cumidino-, monohydrochloride (8CI)  
MF C12 H17 N3 . Cl H



● HCl

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1


L2 10 ANSWERS REGISTRY COPYRIGHT 2002 ACS  
IN 1H-Imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI)  
MF C15 H23 N3



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1


L2 10 ANSWERS REGISTRY COPYRIGHT 2002 ACS  
IN 1H-Indol-5-amine, N-(4,5-dihydro-1H-imidazol-2-yl)-7-(1-methylethyl)- (9CI)  
MF C14 H18 N4  
CI COM



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1


L2 10 ANSWERS REGISTRY COPYRIGHT 2002 ACS  
 IN 1H-Imidazol-2-amine, 4,5-dihydro-N-[2-methyl-6-(1-methylethyl)phenyl]-  
 (9CI)  
 MF C13 H19 N3



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L2 10 ANSWERS REGISTRY COPYRIGHT 2002 ACS  
 IN Benzenamine, N-2-imidazolidinylidene-2,6-bis(1-methylethyl)-,  
 monohydrochloride (9CI)  
 MF C15 H23 N3 . Cl H



● HCl

ALL ANSWERS HAVE BEEN SCANNED

=> fil caplus  
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
140.66	140.87

FULL ESTIMATED COST

FILE 'CAPLUS' ENTERED AT 13:04:22 ON 21 AUG 2002  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 21 Aug 2002 VOL 137 ISS 8  
FILE LAST UPDATED: 20 Aug 2002 (20020820/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

CAS roles have been modified effective December 16, 2001. Please check your SDI profiles to see if they need to be revised. For information on CAS roles, enter HELP ROLES at an arrow prompt or use the CAS Roles thesaurus (/RL field) in this file.

=> s 12  
L3 12 L2

=> d ibib abs hitstr 1-12


L3 ANSWER 1 OF 12 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2001:883550 CAPLUS  
 DOCUMENT NUMBER: 136:320707  
 TITLE: Three-Dimensional common-Feature hypotheses for octopamine agonist 2-(arylimino)imidazolidines  
 AUTHOR(S): Hirashima, Akinori; Morimoto, Masako; Kuvano, Eiichi; Taniguchi, Eiji; Eto, Morifusa  
 CORPORATE SOURCE: Department of Applied Genetics and Pest Management, Kyushu University, Faculty of Agriculture, Graduate School, Fukuoka, Higashi-ku, 812-8581, Japan  
 SOURCE: Bioorganic & Medicinal Chemistry (2001), Volume Date 2002, 10(1), 117-123  
 CODEN: BMECEP; ISSN: 0968-0896  
 PUBLISHER: Elsevier Science Ltd.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB Three-dimensional pharmacophore hypotheses were built from a set of 10 octopamine (OA) agonist 2-(Arylimino)imidazolidines (AIIIs), 2-(Arylimino)thiazolidines (AITs) and 2-(Arylimino)oxazolidines (AIOs). OA agonist activities were detd. using the adenylate cyclase assay in American cockroaches (*P. americana*). Among the 10 common-featured models generated by program Catalyst/HipHop, a hypothesis including a ring arom. (RA), a pos. ionizable (PI) and three hydrophobic aliph. (Hpal) features was considered to be important in evaluating the OA-agonist activity. Active OA agonist 2,6-Et2 AII mapped well onto all the RA, PI and Hpal features of the hypothesis. On the other hand, less active compds. were shown to be difficult to achieve the energetically favorable conformation which is found in the active mols. in order to fit the 3-D common-feature pharmacophore models. Taken together, 2,6-Et2-Ph and foramidine structures are important as OA agonists. The present studies on OA agonists demonstrate that a RA, a PI and three Hpal sites located on the mol. seem to be essential for OA-agonist activity.


IT 63346-74-7 359668-33-0  
 RI: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)  
 (QSAR for octopamine agonist (arylimino)imidazolidines)

RN 63346-74-7 CAPLUS  
 CN 1H-Imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI) (CA INDEX NAME)



RN 359668-33-0 CAPLUS  
 CN 1H-Imidazol-2-amine, 4,5-dihydro-N-[2-methyl-6-(1-methylethyl)phenyl]- (9CI) (CA INDEX NAME)

L3 ANSWER 1 OF 12 CAPLUS COPYRIGHT 2002 ACS (Continued)



REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L3 ANSWER 2 OF 12 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2001:596828 CAPLUS  
 DOCUMENT NUMBER: 135:222824  
 TITLE: Identification of novel inhibitors of calling and in vitro [14C]acetate incorporation by pheromone glands of *Plodia interpunctella*  
 AUTHOR(S): Hirashima, Akinori; Eitaku, Tomohiko; Watanabe, Yasuyuki; Kuvano, Eiichi; Taniguchi, Eiji; Eto, Morifusa  
 CORPORATE SOURCE: Department of Applied Genetics and Pest Management, Faculty of Agriculture, Graduate School, Kyushu University, Fukuoka, 812-8581, Japan  
 SOURCE: Pest Management Science (2001), 57(8), 713-720  
 CODEN: PMSCFQ; ISSN: 1526-498X  
 PUBLISHER: John Wiley & Sons Ltd.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB Some octopamine agonists were found to suppress in vitro biosynthesis of the calling pheromone of the Indian meal moth, *Plodia interpunctella*. Isolated pheromone-gland preps. incorporated sodium [14C]acetate at a linear rate for 3h when incubated with the pheromone biosynthesis activating neuropeptide (PBAN). This incorporation was dependent on the dose of PBAN (up to 0.5 .mu.M). Thin-layer chromatog. of a pheromone-gland ext. revealed quant. incorporation of radioactivity into a product exhibiting the same mobility as (Z,E)-9,12-tetradecadienyl acetate, the main component of the calling pheromone of *P. interpunctella*. Twenty-seven octopamine agonists were initially screened using a calling behavior bioassay of female *P. interpunctella*. Four derivs. with activity in the nanomolar range were identified which were, in order of decreasing pheromonostatic activity: 2-(2,6-diethylphenylimino)thiazolidine > 2-(2,6-dimethylphenylimino)oxazolidine > 2-(2-ethylphenylimino)oxazolidine. These compds. also showed in vitro inhibitory activity in intracellular de novo pheromone biosynthesis. The results of the present study indicate that these derivs. could provide useful information in the characterization and differentiation of octopaminergic receptor types and subtypes.


IT 63346-74-7P 359668-33-0P  
 RI: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (prepn. and pheromonostatic activity of)

RN 63346-74-7 CAPLUS  
 CN 1H-Imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI) (CA INDEX NAME)



RN 359668-33-0 CAPLUS  
 CN 1H-Imidazol-2-amine, 4,5-dihydro-N-[2-methyl-6-(1-methylethyl)phenyl]- (9CI) (CA INDEX NAME)

L3 ANSWER 2 OF 12 CAPLUS COPYRIGHT 2002 ACS (Continued)



REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT





L3 ANSWER 7 OF 12 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1997:594632 CAPLUS

DOCUMENT NUMBER: 127:262678

TITLE:

Preparation of novel indoles and benzothiazoles for cloned human alpha 2 receptors  
 Jeon, Yoon T.; Gluchowski, Charles  
 Synaptic Pharmaceutical Corp., USA  
 PCT Int. Appl., 73 pp.  
 CODEN: PIXXD2

DOCUMENT TYPE:

LANGUAGE:

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9731636	A1	19970904	WO 1997-US3173	19970228
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, HL, HR, NE, SN, TD, TG			
US 5677321	A	19971014	US 1996-608598	19960229
CA 2246813	AA	19970904	CA 1997-2246813	19970228
AU 9720604	A1	19970916	AU 1997-20604	19970228
AU 704439	B2	19990422		
EP 900080	A1	19990310	EP 1997-908782	19970228
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, HC, PT, IE, FI			
JP 2000506144	T2	20000523	JP 1997-531156	19970228
US 5948804	A	19990907	US 1997-926316	19970905
US 6040451	A	20000321	US 1999-345470	19990630
US 6159998	A	20001212	US 2000-492505	20000127
US 6303643	B1	20011016	US 2000-690620	20000107
US 2002049239	A1	20020425	US 2001-965944	20010928
PRIORITY APPLN. INFO.:			US 1996-608598	A 19960229
			WO 1997-US3173	W 19970228
			US 1997-926316	A1 19970905
			US 1999-345470	A1 19990630
			US 2000-492505	A1 20000127
			US 2000-690620	A1 20000107

OTHER SOURCE(S):  
GI

MARPAT 127:262678

\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB The title compds. [I-IV; R1-R3 = H, C1-7 alkyl, C2-7 alkenyl, alkynyl; R4-R6 = H, halo, Oh, etc.; R7 = H, NH, C1-7 alkyl, etc.; R8 = H, C1-7 alkyl, C2-7 alkenyl, etc.; R9 = H, Ph, C1-7 alkyl, etc.; X = CH2, O, NH, S] which are selective for cloned human alpha 2 receptors and therefore

L3 ANSWER 7 OF 12 CAPLUS COPYRIGHT 2002 ACS (Continued)


HO<sub>2</sub>C-CH<sub>2</sub>-CH<sub>2</sub>-CO<sub>2</sub>H

L3 ANSWER 7 OF 12 CAPLUS COPYRIGHT 2002 ACS (Continued)

useful for lowering intraocular pressure, for treating presbyopia, migraine, hypertension, alc. withdrawal, drug addiction, rheumatoid arthritis, ischemic pain, spasticity, diarrhea, nasal congestion, urinary incontinence as well as for use as analgesics, sedatives, anesthetics, cognition enhancers and ocular vasoconstriction agents, were prepd. Thus, reaction of 7-bromo-5-aminoindole with 2-imidazoline-2-sulfonic acid (ISA) afforded 46a: [R1-R5 = H; R6 = Br; R7-R9 = H; X = N] which showed pEC<sub>50</sub> of 9.36 at alpha 2 receptor.

IT 196204-74-7P 196204-75-8P  
 RI: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (prepn. of novel indoles and benzothiazoles for cloned human alpha 2 receptors)


RN 196204-74-7 CAPLUS  
 CN 1H-Indol-5-amine, N-(4,5-dihydro-1H-imidazol-2-yl)-7-(1-methylethyl)- (9CI) (CA INDEX NAME)



RN 196204-75-8 CAPLUS  
 CN Butanedioic acid, compd. with N-(4,5-dihydro-1H-imidazol-2-yl)-7-(1-methylethyl)-1H-indol-5-amine (1:1) (9CI) (CA INDEX NAME)

CH 1

CRN 196204-74-7  
 CMF C14 H18 N4



CH 2

CRN 110-15-6  
 CMF C4 H6 O4

L3 ANSWER 8 OF 12 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1995:303934 CAPLUS

DOCUMENT NUMBER: 122:77274

TITLE: Pharmacology of the octopamine receptor from locust central nervous tissue (OAR3)

AUTHOR(S): Roeder, Thomas  
 CORPORATE SOURCE: Zool. Inst., Univ. Hamburg, Hamburg, D-20146, Germany  
 SOURCE: British Journal of Pharmacology (1995), 114(1), 210-16  
 CODEN: BJPCBM; ISSN: 0007-1188

PUBLISHER: Stockton


DOCUMENT TYPE: Journal

LANGUAGE: English

AB The present study characterized highly effective agonists from different classes of compds. for the neuronal octopamine receptor (OAR3) of the migratory locust (*Locusta migratoria* L.). Biogenic amines and phenyliminoimidazolidines (PIIs) were employed for the study of structure-activity relationships. The highest affinity PIIs were predominantly those with the substitutions at the positions 2 and 4 of the phenolic ring (e.g. NC 7, K<sub>i</sub> = 0.3 nM, NC8, K<sub>i</sub> = 0.81 nM). Substitutions at these positions always had pos. effects on the affinity of the resp. agonists. Substitutions at the positions 3, 5 and 6, however, always had neg. effects on the affinity. At the position one of the phenolic ring, heterocyclic substituents are preferred. Some PIIs had a more than 30 times higher affinity for OARs than for .alpha.-adrenoceptors which are the vertebrate homologues of the insect octopamine receptors. The only non-PII with subnanomolar affinity was the aminooxazoline deriv. AC 6 (K<sub>i</sub> = 0.92 nM). A variety of substances with known insecticidal activity such as chlordimeform, demethylchlordimeform, amitraz or AC 6 had high affinity for the locust neuronal octopamine receptor.

IT RI: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)  
 (structure-activity relationship of agonists for locust neuronal octopamine receptor)

RN 63346-74-7 CAPLUS  
 CN 1H-Imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI) (CA INDEX NAME)




L3 ANSWER 9 OF 12 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1989:95240 CAPLUS  
 DOCUMENT NUMBER: 110:95240  
 TITLE: Preparation of 2-(phenylimino)imidazolidines as .alpha.1-adrenergic agonists  
 INVENTOR(S): Esser, Franz; Staehle, Helmut; Koeppel, Herbert; Speck, Georg; Miesau, Joachim; Pichler, Ludwig; Lehr, Erich  
 PATENT ASSIGNEE(S): Boehringer Ingelheim K.-G., Fed. Rep. Ger.  
 SOURCE: Ger. Offen., 7 pp.  
 CODEN: GWXXBX  
 DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 3712385	A1	19881027	DE 1987-3712385	19870411

OTHER SOURCE(S): CASREACT 110:95240; MARPAT 110:95240  
 GI




AB The title compds. [I; R1, R2 = F, Cl, Br, iodo; R3 = (substituted) C1-4 alkyl] and pharmaceutically acceptable salts were prep'd. as CNS agents and cyto- and cardioprotectants. KSCN in acetone was treated with PhCOCl at 15.degree. and 2-chloro-4-isopropylaniline was added. The mixt. was refluxed 3.25 h to give 70.5% (2-chloro-4-isopropylphenyl)thiourea. The latter was sequentially refluxed with MeI in MeOH, refluxed with H2NCH2CH2NH2 in MeOH, stirred with 5N NaOH, and treated with Br in CHCl3 at 0-8.degree. to give 2-(2-chloro-4-isopropylphenylimino)imidazolidine.HBr. The latter at 1 mg/kg in mice increased survival in a hypoxia screen from 40% (controls) to 70%.

IT 118955-15-0P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of, as CNS agent and cardio- and cytoprotectant)


RN 118955-15-0 CAPLUS  
 CN 1H-Imidazol-2-amine, N-[2-bromo-6-chloro-4-(1-methylethyl)phenyl]-4,5-dihydro-, monohydrobromide (9CI) (CA INDEX NAME)

L3 ANSWER 9 OF 12 CAPLUS COPYRIGHT 2002 ACS (Continued)



● HBr

IT 118954-98-1P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of, as CNS agent, cyto- and cardioprotectant)  
 RN 118954-98-1 CAPLUS  
 CN 1H-Imidazol-2-amine, N-[2-bromo-6-chloro-4-(1-methylethyl)phenyl]-4,5-dihydro- (9CI) (CA INDEX NAME)




L3 ANSWER 10 OF 12 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1985:593431 CAPLUS  
 DOCUMENT NUMBER: 103:193431  
 TITLE: Phenyliminoimidazolidines. Characterization of a class of potent agonists of octopamine-sensitive adenylate cyclase and their use in understanding the pharmacology of octopamine receptors  
 AUTHOR(S): Nathanson, James A.  
 CORPORATE SOURCE: Dep. Neurol., Harvard Med. Sch., Boston, MA, 02114, USA  
 SOURCE: Mol. Pharmacol. (1985), 28(3), 254-68  
 CODEN: MOPMAJ; ISSN: 0026-195X  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB Approx. 30 substituted phenyliminoimidazolidines (PII) were exam'd. for agonist and antagonist effects on the highly enriched and specific octopamine (O)-sensitive adenylate cyclase (AC) present in the firefly light organ, as well as on ACs present in other invertebrate and some had vertebrate tissues. Several derivs. were extremely active and some had potencies exceeding those of any previously described agonists of O-sensitive AC. Stimulation by the potent PIIs was reversible, nonadditive to that caused by O, and could be antagonized by antagonists such as cyproheptadine, phentolamine, and propranolol. The inhibitory consts. agreed well with those for inhibiting O stimulation. Certain PII derivs. acted as partial agonists and some as antagonists of O stimulation. Structure-activity relationships revealed, among other things, that short-chain alkyl substitution in the 2- and 6-Phe positions enhanced activity, as did further substitution of 4-halo, 4-Me, or 4-hydroxy substituents. 4-Amino or N-alkyl substitution decreased activity. Structurally related benzylimidazoline derivs. such as tolazoline and naphazoline were partial O agonists, generally less active than the PIIs. Comparison, in 3 invertebrate species, of the effects of the PIIs and 2 other chem. classes of O agonists demonstrated clearcut differences in species responsiveness. Other comparative studies revealed that the agonist activity of the potent PIIs was specific for tissues containing an O-sensitive AC; ACs activated by dopamine or by .beta.1- or .beta.2-adrenergic agonists were unaffected by these compds. The active PIIs affected a class of O receptors distinct from mammalian .alpha.1- or .alpha.2-adrenergic receptors. These O receptors also appeared distinct from mammalian 5-HT1 and 5-HT2 receptors. Correlative physiol. studies in insects revealed that the active PIIs mimicked O and were potent activators of light emission in the firefly light organ. The PIIs also cause disruption of motor and feeding behavior in tobacco hornworms, leading to insect death, an effect which was markedly potentiated by phosphodiesterase inhibitors.


IT 63346-74-7  
 RL: BIOL (Biological study)  
 (octopamine-sensitive adenylate cyclase activation by, in light organ of firefly, structure in relation to)

RN 63346-74-7 CAPLUS  
 CN 1H-Imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI) (CA INDEX NAME)


L3 ANSWER 10 OF 12 CAPLUS COPYRIGHT 2002 ACS (Continued)




L3 ANSWER 11 OF 12 CAPLUS COPYRIGHT 2002 ACS  
 ACCESSION NUMBER: 1977:447931 CAPLUS  
 DOCUMENT NUMBER: 87:47931  
 TITLE: Structure-activity relations and problems related to the mechanism of action of clonidine  
 AUTHOR(S): Rouot, Bruno; Leclerc, Gerard; Vermuth, Camille-Georges; Miesch, Francois; Schwartz, Jean  
 CORPORATE SOURCE: Fac. Pharm., Univ. Louis-Pasteur, Strasbourg, Fr.  
 SOURCE: J. Pharmacol. (1977), 8(1), 95-106  
 CODEN: JNPBAG  
 DOCUMENT TYPE: Journal  
 LANGUAGE: French  
 GI




AB All 26 clonidine analogs (I) studied showed peripheral .alpha.-sympathomimetic activity, with IPS 56 (I; R = 2,3-dichloro) [15327-44-3] having the greatest hypertensive effect in demedullated rats. The results correlated with Es (steric const.) and F (sum of the field effect of the substituents) of the Hansch equation (1971). No such correlation was obsd. for hypotensive activity in intact rats. Compds. which had hypotensive activity also had high .alpha.-sympathomimetic activity, local anesthetic activity (rabbit cornea), and similar lipophilicity. However, the results were not sufficient to conclude that .alpha.-adrenergic mechanisms are involved in the hypotensive effects of clonidine and its analogs.  
 IT 63346-74-7  
 RL: BAC (Biological activity or effector, except adverse); BIOL (Biological study)  
 (hypotensive and .alpha.-sympathomimetic activity of)  
 RN 63346-74-7 CAPLUS  
 CN 1H-imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI) (CA INDEX NAME)



L3 ANSWER 12 OF 12 CAPLUS COPYRIGHT 2002 ACS  
 ACCESSION NUMBER: 1976:456532 CAPLUS  
 DOCUMENT NUMBER: 85:56532  
 TITLE: Clonidine and related analogues. Quantitative correlations  
 AUTHOR(S): Rouot, Bruno; Leclerc, Gerard; Vermuth, Camille G.; Miesch, Francois; Schwartz, Jean  
 CORPORATE SOURCE: Fac. Pharm., Univ. Louis Pasteur, Strasbourg, Fr.  
 SOURCE: J. Med. Chem. (1976), 19(8), 1049-54  
 CODEN: JMCHAR  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 GI



AB A series of 22 derivs. of clonidine-HCl (I) [4205-91-8] were prepd. by the cyclization reaction of ethylenediamine with an S-methylisothiuronium salt deriv. and the main physicochem. parameters (log P, .DELTA.NM, pKa) detd. Quant. correlations between peripheral .alpha.-mimetic action (pithed rats) and physicochem. parameters pointed out the crit. role of the steric effect of ortho substituents. Attempted quant. correlations between physicochem. parameters and central hypotensive activity were unsuccessful. The mechanism of action of I is discussed.  
 IT 59465-43-99  
 RL: SPN (Synthetic preparation); PREP (Preparation) (prepn. and blood pressure response to)  
 RN 59465-43-9 CAPLUS  
 CN Benzenamine, N-2-imidazolidinylidene-2,6-bis(1-methylethyl)-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

=> log y

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

53.06

193.93

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-7.43

-7.43

STN INTERNATIONAL LOGOFF AT 13:05:10 ON 21 AUG 2002

Welcome to STN International! Enter x:x

LOGINID:sssptal600rxa

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Apr 08 "Ask CAS" for self-help around the clock  
NEWS 3 Apr 09 BEILSTEIN: Reload and Implementation of a New Subject Area  
NEWS 4 Apr 09 ZDB will be removed from STN  
NEWS 5 Apr 19 US Patent Applications available in IFICDB, IFIPAT, and IFIUDB  
NEWS 6 Apr 22 Records from IP.com available in CAPLUS, HCAPLUS, and ZCAPLUS  
NEWS 7 Apr 22 BIOSIS Gene Names now available in TOXCENTER  
NEWS 8 Apr 22 Federal Research in Progress (FEDRIP) now available  
NEWS 9 Jun 03 New e-mail delivery for search results now available  
NEWS 10 Jun 10 MEDLINE Reload  
NEWS 11 Jun 10 PCTFULL has been reloaded  
NEWS 12 Jul 02 FOREGE no longer contains STANDARDS file segment  
NEWS 13 Jul 22 USAN to be reloaded July 28, 2002;  
saved answer sets no longer valid  
NEWS 14 Jul 29 Enhanced polymer searching in REGISTRY  
NEWS 15 Jul 30 NETFIRST to be removed from STN  
NEWS 16 Aug 08 CANCERLIT reload  
NEWS 17 Aug 08 PHARMAMarketLetter(PHARMAML) - new on STN  
NEWS 18 Aug 08 NTIS has been reloaded and enhanced  
NEWS 19 Aug 09 JAPIO to be reloaded August 25, 2002  
NEWS 20 Aug 19 Aquatic Toxicity Information Retrieval (AQUIRE)  
now available on STN  
NEWS 21 Aug 19 IFIPAT, IFICDB, and IFIUDB have been reloaded  
NEWS 22 Aug 19 The MEDLINE file segment of TOXCENTER has been reloaded  
  
NEWS EXPRESS February 1 CURRENT WINDOWS VERSION IS V6.0d,  
CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP),  
AND CURRENT DISCOVER FILE IS DATED 05 FEBRUARY 2002  
  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 12:12:10 ON 21 AUG 2002

=> fil reg		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'REGISTRY' ENTERED AT 12:12:22 ON 21 AUG 2002  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 American Chemical Society (ACS)

STRUCTURE FILE UPDATES: 20 AUG 2002 HIGHEST RN 444542-98-7  
DICTIONARY FILE UPDATES: 20 AUG 2002 HIGHEST RN 444542-98-7

TSCA INFORMATION NOW CURRENT THROUGH MAY 20, 2002

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.


Crossover limits have been increased. See HELP CROSSOVER for details.

Calculated physical property data is now available. See HELP PROPERTIES  
for more information. See STNote 27, Searching Properties in the CAS  
Registry File, for complete details:  
<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

=>  
Uploading 10058456.str

L1 STRUCTURE UPLOADED

=> d  
L1 HAS NO ANSWERS  
L1 STR



Structure attributes must be viewed using STN Express query preparation.

=> s l1  
SAMPLE SEARCH INITIATED 12:12:36 FILE 'REGISTRY'  
SAMPLE SCREEN SEARCH COMPLETED - 990 TO ITERATE

100.0% PROCESSED 990 ITERATIONS  
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)  
SEARCH TIME: 00.00.01

50 ANSWERS

Examiner Anderson 703-605-1157

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*  
 BATCH \*\*COMPLETE\*\*  
 PROJECTED ITERATIONS: 17913 TO 21687  
 PROJECTED ANSWERS: 4943 TO 7017

L2 50 SEA SSS SAM L1

=> s l1 full  
 FULL SEARCH INITIATED 12:12:40 FILE 'REGISTRY'  
 FULL SCREEN SEARCH COMPLETED - 20061 TO ITERATE

100.0% PROCESSED 20061 ITERATIONS 5848 ANSWERS  
 SEARCH TIME: 00.00.02

L3 5848 SEA SSS FUL L1

=> fil caplus		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	140.28	140.49

FILE 'CAPLUS' ENTERED AT 12:12:46 ON 21 AUG 2002  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 21 Aug 2002 VOL 137 ISS 8  
 FILE LAST UPDATED: 20 Aug 2002 (20020820/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

CAS roles have been modified effective December 16, 2001. Please check your SDI profiles to see if they need to be revised. For information on CAS roles, enter HELP ROLES at an arrow prompt or use the CAS Roles thesaurus (/RL field) in this file.

=> s l3  
 L4 9033 L3

=> fil reg		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.40	140.89

FILE 'REGISTRY' ENTERED AT 12:12:55 ON 21 AUG 2002  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

Examiner Anderson 703-605-1157



PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 American Chemical Society (ACS)

STRUCTURE FILE UPDATES: 20 AUG 2002 HIGHEST RN 444542-98-7  
DICTIONARY FILE UPDATES: 20 AUG 2002 HIGHEST RN 444542-98-7

TSCA INFORMATION NOW CURRENT THROUGH MAY 20, 2002

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Calculated physical property data is now available. See HELP PROPERTIES  
for more information. See STNote 27, Searching Properties in the CAS  
Registry File, for complete details:  
<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

=>


Uploading 10058456.str

L5 STRUCTURE UPLOADED

=> d

L5 HAS NO ANSWERS

L5 STR



G1 H,X,O,C

G2 C,H,X

Structure attributes must be viewed using STN Express query preparation.

=> s 15 subset=13 full

FULL SUBSET SEARCH INITIATED 12:15:46 FILE 'REGISTRY'

FULL SUBSET SCREEN SEARCH COMPLETED - 5848 TO ITERATE

100.0% PROCESSED 5848 ITERATIONS  
SEARCH TIME: 00.00.01

3794 ANSWERS

L6 3794 SEA SUB=L3 SSS FUL L5

=>

Uploading 10058456.str


Examiner Anderson 703-605-1157

L7 STRUCTURE UPLOADED

=> d

L7 HAS NO ANSWERS

L7 STR



G1 H,X,O,C

G2 C,H,X

Structure attributes must be viewed using STN Express query preparation.

=> s l7 subset=l5 full

L5 MAY NOT BE USED HERE

The L-number must have been created by a search in this file. To see all L-numbers defined in this session, enter DISPLAY HISTORY at an arrow prompt (=). For additional information on subset searching in this file, enter HELP SUBSET.

ENTER SUBSET L# OR (END):6

6 IS NOT A VALID L#

L-numbers must be in the range L1-L999.

ENTER SUBSET L# OR (END):16

FULL SUBSET SEARCH INITIATED 12:17:26 FILE 'REGISTRY'

FULL SUBSET SCREEN SEARCH COMPLETED - 3794 TO ITERATE

100.0% PROCESSED 3794 ITERATIONS  
SEARCH TIME: 00.00.01

1495 ANSWERS

L8 1495 SEA SUB=L6 SSS FUL L7

=> fil caplus

COST IN U.S. DOLLARS

SINCE FILE

ENTRY

69.14

TOTAL

SESSION

210.03

FULL ESTIMATED COST

FILE 'CAPLUS' ENTERED AT 12:17:37 ON 21 AUG 2002

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is

Examiner Anderson 703-605-1157

held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 21 Aug 2002 VOL 137 ISS 8  
FILE LAST UPDATED: 20 Aug 2002 (20020820/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

CAS roles have been modified effective December 16, 2001. Please check your SDI profiles to see if they need to be revised. For information on CAS roles, enter HELP ROLES at an arrow prompt or use the CAS Roles thesaurus (/RL field) in this file.

=> s l8  
L9 7548 L8

=> fil reg	SINCE FILE	TOTAL
COST IN U.S. DOLLARS	ENTRY	SESSION
FULL ESTIMATED COST	1.19	211.22

FILE 'REGISTRY' ENTERED AT 12:19:37 ON 21 AUG 2002  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 American Chemical Society (ACS)

STRUCTURE FILE UPDATES: 20 AUG 2002 HIGHEST RN 444542-98-7  
DICTIONARY FILE UPDATES: 20 AUG 2002 HIGHEST RN 444542-98-7

TSCA INFORMATION NOW CURRENT THROUGH MAY 20, 2002

Please note that search-term pricing does apply when conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.


Calculated physical property data is now available. See HELP PROPERTIES for more information. See STNote 27, Searching Properties in the CAS Registry File, for complete details:  
<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

=>  
Uploading 10058456.str

L10 STRUCTURE UPLOADED

=> d  
L10 HAS NO ANSWERS  
L10 STR

Examiner Anderson 703-605-1157



G1 H,X,O,C

G2 C,H,X

Structure attributes must be viewed using STN Express query preparation.

=> s l10 subset=l8 full

FULL SUBSET SEARCH INITIATED 12:20:03 FILE 'REGISTRY'  
FULL SUBSET SCREEN SEARCH COMPLETED - 1495 TO ITERATE

100.0% PROCESSED 1495 ITERATIONS  
SEARCH TIME: 00.00.01

218 ANSWERS

L11 218 SEA SUB=L8 SSS FUL L10

=> fil caplus  
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
33.43	244.65

FULL ESTIMATED COST

FILE 'CAPLUS' ENTERED AT 12:20:07 ON 21 AUG 2002  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 21 Aug 2002 VOL 137 ISS 8  
FILE LAST UPDATED: 20 Aug 2002 (20020820/ED)

This file contains CAS Registry Numbers for easy and accurate

Examiner Anderson 703-605-1157

substance identification.

CAS roles have been modified effective December 16, 2001. Please check your SDI profiles to see if they need to be revised. For information on CAS roles, enter HELP ROLES at an arrow prompt or use the CAS Roles thesaurus (/RL field) in this file.

=> s l11

L12            328 L11

=> s l12 and urinary

107190 URINARY

L13

2 L12 AND URINARY

=> d ibib abs hitstr 1-2

L13 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1997:318325 CAPLUS

DOCUMENT NUMBER: 127:13229

TITLE: The peripheral action of clonidine analog ST-91: involvement of atrial natriuretic factor

AUTHOR(S): Gutkowska, Jolanta; Mukaddam-Daheer, Suhayla; Tremblay, Johanne

CORPORATE SOURCE: Laboratory Cardiovascular Biochemistry, Centre Recherche Hotel-Dieu Montreal, Universite Montreal, Montreal, QC, H2W 1T8, Can.

SOURCE: Journal of Pharmacology and Experimental Therapeutics (1997), 281(2), 670-676

CODEN: JPETAB; ISSN: 0022-3565

PUBLISHER: Williams &amp; Wilkins

DOCUMENT TYPE: Journal


LANGUAGE: English

AB It is generally thought that the cardiovascular and renal effects of clonidine, an alpha-2 adrenergic agonist, are mediated by central mechanisms. Our previous work has shown that diuresis and natriuresis caused by central administration of clonidine are mediated by an enhanced release of atrial natriuretic factor (ANF). Because clonidine has been shown to have peripheral actions the objective of the present study was to det. whether ANF is also involved in these actions. Studies were performed with use of a structural clonidine analog, ST-91, which does not cross the blood-brain barrier. I.v. injection of various doses (0-250 .mu.g/rat) of ST-91 into conscious, normally hydrated female Sprague-Dawley rats (200-250 g) produced dose-related increases in urinary output, which were accompanied by significant increases in urinary sodium, potassium and cGMP excretion. Compared with saline, the highest dose of ST-91 (250 .mu.g/rat) during the first hour of treatment significantly ( $P < .001$ ,  $n = 18$ ) enhanced urinary output (0.2 +/- 0.1 vs. 3.0 +/- 1.1 mL/h) and excretion of sodium (28 +/- 4 vs. 345 +/- 50 .mu.mol/h), potassium (10 +/- 4 vs. 165 +/- 37 .mu.mol/h) and cGMP (191 +/- 29 vs. 1340 +/- 322 pmol/h). The biol. marker of ANF. These renal responses were assoc. with increased plasma ANF (59 +/- 28 pg/mL,  $P < .001$ ,  $n = 12$ ), measured 10 min after ST-91 (250 .mu.g/rat), which remained elevated for at least 1 h ( $P < .01$ ,  $n = 6$ ). The enhanced renal responses that were induced by 10 .mu.g ST-91 were partially, yet significantly inhibited by yohimbine (50 .mu.g), an alpha-2 antagonist. On the other hand, efaroxan (500 .mu.g), an 11 imidazoline receptor antagonist, showed a stronger inhibitory effect, whereas naloxone (0.8 mg) had no effect. Pretreatment of rats with anti-ANF reduced the diuretic and natriuretic effects of ST-91. These results indicate that the renal effects of ST-91 are mediated by imidazoline as well as by alpha-2 adrenergic receptors, but not by opioid receptors. Furthermore, the renal effects evoked by ST-91 are mediated by ANF.

IT 4749-61-5, ST-91  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(peripheral action of clonidine analog ST-91: involvement of atrial natriuretic factor)

RN 4749-61-5 CAPLUS  
 CN 1H-imidazol-2-amine, N-(2,6-diethylphenyl)-4,5-dihydro-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

L13 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1990:1110 CAPLUS

DOCUMENT NUMBER: 112:1110

TITLE: .alpha.2-Adrenergic receptors and the sodium/hydrogen ion exchanger in the intestinal epithelial cell line, HT-29

AUTHOR(S): Cantilello, Horacio F.; Lanier, Stephen M.

CORPORATE SOURCE: Massachusetts Gen. Hosp., Harvard Med. Sch., Boston, MA, 02114, USA

SOURCE: J. Biol. Chem. (1989), 264(27), 16000-7

CODEN: JBCHA3; ISSN: 0021-9258


DOCUMENT TYPE: Journal

LANGUAGE: English

AB The effect of .alpha.2-adrenergic receptors (.alpha.2-AR) activation on basal and stimulated Na<sup>+</sup>/H<sup>+</sup> exchange was studied in epithelial cells isolated from human colon (HT-29 adenocarcinoma cells). Na<sup>+</sup>/H<sup>+</sup> exchange was measured by quantitation of intracellular H<sup>+</sup> ion concn. (acetoxymethyl ester 2,7-bis(carboxyethyl)-5(6)-carboxyfluorescein) and 22Na<sup>+</sup> uptake. HT-29 cells expressed an amiloride-sensitive Na<sup>+</sup>/H<sup>+</sup> exchanger that was activated by redn. of intracellular pH (pHi) to 6.0 but was quiescent at a physiol. pHi. The rapid alkalization obsd. after acid loading (0.57 pH units/min/104 cells) was dependent on external Na and was blocked by amiloride (Ki .apprx. 2.1 .mu.M). .alpha.2-AR agonists clonidine and UK-14304 inhibited forskolin-activated adenylyl cyclase, these compds. did not alter basal Na<sup>+</sup>/H<sup>+</sup> exchange. Stimulated Na<sup>+</sup>/H<sup>+</sup> exchange was similarly unaffected by epinephrine. In contrast, stimulated Na<sup>+</sup>/H<sup>+</sup> exchanger activity was completely inhibited by the selective .alpha.2-agonists clonidine, UK-14304, and guanabenz. This inhibitory effect was not blocked by the .alpha.2-AR antagonist rauwolfscine, and it is likely due to a direct interaction with the exchanger mol. itself. Structure/activity studies indicated that the compds. inhibiting exchanger activity possess either an imidazoline or guanidinium moiety. Although these mols. bear structural similarity to amiloride, they did not inhibit the amiloride-sensitive epithelial Na channel in toad urinary bladder, suggesting that these compds. may be useful as amiloride-like ligands selective for the Na<sup>+</sup>/H<sup>+</sup> exchanger. Evidently, in the HT-29 intestinal cell line, in contrast to observations in other tissues, .alpha.2-AR are not coupled to the Na<sup>+</sup>/H<sup>+</sup> exchanger, suggesting that the cell-signaling mechanisms utilized by the .alpha.2-AR are tissue specific.

IT 4749-61-5, ST-91  
 RL: BIOL (Biological study)  
 (hydrogen ion-sodium exchange transport inhibition by, in intestine epithelial cells of human, structure in relation to)

RN 4749-61-5 CAPLUS  
 CN 1H-imidazol-2-amine, N-(2,6-diethylphenyl)-4,5-dihydro-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

Examiner Anderson 703-605-1157

=> FIL STNGUIDE	SINCE FILE	TOTAL
COST IN U.S. DOLLARS	ENTRY	SESSION
	11.59	256.24
FULL ESTIMATED COST		
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
	-1.24	-1.24
CA SUBSCRIBER PRICE		

FILE 'STNGUIDE' ENTERED AT 12:21:56 ON 21 AUG 2002  
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT  
 COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE  
 AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.  
 LAST RELOADED: Aug 16, 2002 (20020816/UP).

=> fil reg	SINCE FILE	TOTAL
COST IN U.S. DOLLARS	ENTRY	SESSION
	0.06	256.30
FULL ESTIMATED COST		
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
	0.00	-1.24
CA SUBSCRIBER PRICE		

FILE 'REGISTRY' ENTERED AT 12:22:21 ON 21 AUG 2002  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2002 American Chemical Society (ACS)

STRUCTURE FILE UPDATES: 20 AUG 2002 HIGHEST RN 444542-98-7  
 DICTIONARY FILE UPDATES: 20 AUG 2002 HIGHEST RN 444542-98-7

TSCA INFORMATION NOW CURRENT THROUGH MAY 20, 2002

Please note that search-term pricing does apply when  
 conducting SmartSELECT searches.


Crossover limits have been increased. See HELP CROSSOVER for details.

Calculated physical property data is now available. See HELP PROPERTIES  
 for more information. See STNote 27, Searching Properties in the CAS  
 Registry File, for complete details:  
<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

=>  
 Uploading 10058456.str

L14 STRUCTURE UPLOADED

=> d  
 L14 HAS NO ANSWERS  
 L14 STR



G1 H,X,O,C

G2 C,H,X

Structure attributes must be viewed using STN Express query preparation.

=> s 114 subset=18 full  
FULL SUBSET SEARCH INITIATED 12:23:38 FILE 'REGISTRY'  
FULL SUBSET SCREEN SEARCH COMPLETED - 1495 TO ITERATE

100.0% PROCESSED 1495 ITERATIONS  
SEARCH TIME: 00.00.01


4 ANSWERS

L15 4 SEA SUB=L8 SSS FUL L14

=> d scan




L15 4 ANSWERS REGISTRY COPYRIGHT 2002 ACS  
IN 1H-Imidazol-2-amine, 4,5-dihydro-N-[2-methyl-6-(1-methylethyl)phenyl]-  
(9CI)  
MF C13 H19 N3



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1


L15 4 ANSWERS REGISTRY COPYRIGHT 2002 ACS  
IN Benzenamine, N-2-imidazolidinylidene-2,6-bis(1-methylethyl)-,  
monohydrochloride (9CI)  
MF C15 H23 N3 . Cl H



● HCl

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1


L15 4 ANSWERS REGISTRY COPYRIGHT 2002 ACS  
IN 1H-Imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI)  
MF C15 H23 N3



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L15 4 ANSWERS REGISTRY COPYRIGHT 2002 ACS  
IN 1H-Imidazol-2-amine, 4,5-dihydro-N-hydroxy-N-[3-(1-methylethyl)phenyl]-,  
monohydrochloride (9CI)  
MF C12 H17 N3 O . Cl H



● HCl

ALL ANSWERS HAVE BEEN SCANNED

=> fil caplus  
 COST IN U.S. DOLLARS  
 FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
34.19	290.49

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)  
 CA SUBSCRIBER PRICE

SINCE FILE	TOTAL
ENTRY	SESSION
0.00	-1.24

FILE 'CAPLUS' ENTERED AT 12:23:57 ON 21 AUG 2002  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 21 Aug 2002 VOL 137 ISS 8  
 FILE LAST UPDATED: 20 Aug 2002 (20020820/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

CAS roles have been modified effective December 16, 2001. Please check your SDI profiles to see if they need to be revised. For information on CAS roles, enter HELP ROLES at an arrow prompt or use the CAS Roles thesaurus (/RL field) in this file.

=> s l15  
 L16 10 L15  
 => d ibib abs hitstr 1-10

L16 ANSWER 1 OF 10 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2001:88350 CAPLUS

DOCUMENT NUMBER: 136:320707

TITLE:

AUTHOR(S):

CORPORATE SOURCE:

SOURCE:

PUBLISHER:

DOCUMENT TYPE:


LANGUAGE:

AB Three-dimensional pharmacophore hypotheses were built from a set of 10 octopamine (OA) agonist 2-(Arylimino)imidazolidines (AIIIs), 2-(Arylimino)thiazolidines (AITs) and 2-(Arylimino)oxazolidines (AIOs). OA agonist activities were detd. using the adenylate cyclase assay in American cockroaches (*P. americana*). Among the 10 common-featured models generated by program Catalyst/HipHop, a hypothesis including a ring arom. (RA), a pos. ionizable (PI) and three hydrophobic aliph. (HpAl) features was considered to be important in evaluating the OA-agonist activity. Active OA agonist 2,6-Et2 AII mapped well onto all the RA, PI and HpAl features of the hypothesis. On the other hand, less active compds. were shown to be difficult to achieve the energetically favorable conformation which is found in the active mols. in order to fit the 3-D common-feature pharmacophore models. Taken together, 2,6-Et2-Ph and foramidine structures are important as OA agonists. The present studies on OA agonists demonstrate that a RA, a PI and three HpAl sites located on the mol. seem to be essential for OA-agonist activity.

IT 63346-74-7 359668-33-0  
 RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)  
 (QSAR for octopamine agonist (arylimino)imidazolidines)

RN 63346-74-7 CAPLUS


CN 1H-Imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI)  
 (CA INDEX NAME)



RN 359668-33-0 CAPLUS

CN 1H-Imidazol-2-amine, 4,5-dihydro-N-[2-methyl-6-(1-methylethyl)phenyl]- (9CI) (CA INDEX NAME)

L16 ANSWER 1 OF 10 CAPLUS COPYRIGHT 2002 ACS (Continued)



REFERENCE COUNT:

17

THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L16 ANSWER 2 OF 10 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2001:596828 CAPLUS

DOCUMENT NUMBER: 135:222824

TITLE:

AUTHOR(S):

CORPORATE SOURCE:

SOURCE:

PUBLISHER:

DOCUMENT TYPE:


LANGUAGE:

AB Some octopamine agonists were found to suppress in vitro biosynthesis of the calling pheromone of the Indian meal moth, *Plodia interpunctella*. Isolated pheromone-gland preps. incorporated sodium [14C]acetate at a linear rate for 3h when incubated with the pheromone biosynthesis activating neuropeptide (PBAN). This incorporation was dependent on the dose of PBAN (up to 0.5 .mu.M). Thin-layer chromatog. of a pheromone-gland ext. revealed quant. incorporation of radioactivity into a product exhibiting the same mobility as (Z,E)-9,12-tetradecadienyl acetate, the main component of the calling pheromone of *P. interpunctella*. Twenty-seven octopamine agonists were initially screened using a calling behavior bioassay of female *P. interpunctella*. Four derivs. with activity in the nanomolar range were identified which were, in order of decreasing pheromonostatic activity: 2-(2,6-diethylphenylimino)thiazolidine > 2-(2,6-dimethylphenylimino)oxazolidine > 2-(2-ethylphenylimino)oxazolidine. These compds. also showed in vitro inhibitory activity in intracellular de novo pheromone biosynthesis. The results of the present study indicate that these derivs. could provide useful information in the characterization and differentiation of octopaminergic receptor types and subtypes.

IT 63346-74-7P 359668-33-0P  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (prep. and pheromonostatic activity of)

RN 63346-74-7 CAPLUS


CN 1H-Imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI)  
 (CA INDEX NAME)



RN 359668-33-0 CAPLUS

CN 1H-Imidazol-2-amine, 4,5-dihydro-N-[2-methyl-6-(1-methylethyl)phenyl]- (9CI) (CA INDEX NAME)

L16 ANSWER 2 OF 10 CAPLUS COPYRIGHT 2002 ACS (Continued)




REFERENCE COUNT:

41

THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L16 ANSWER 3 OF 10 CAPLUS COPYRIGHT 2002 ACS  
 ACCESSION NUMBER: 2001:37784 CAPLUS  
 DOCUMENT NUMBER: 134:231513  
 TITLE: Synthesis, structure, and binding of some 2-imidazolines to rat brain  $\alpha$ -1 and  $\alpha$ -2-adrenergic receptors  
 AUTHOR(S): Sazewski, F.; Kobierska, E.; Debowski, T.; Charakchiwa-Minol, S.; Mokrosz, M.; Gdaniec, M.; Nowak, E.  
 CORPORATE SOURCE: Department of Chemical Technology of Drug and Organic Chemistry, Medical University of Gdansk, Pol.  
 SOURCE: Archiv der Pharmazie (Weinheim, Germany) (2000), 333(12), 425-430  
 CODEN: ARPMAS; ISSN: 0365-6233  
 PUBLISHER: Wiley-VCH Verlag GmbH  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 134:231513  
 AB A series of novel 2-[(2-aminophenyl)imino]imidazolinium salts and N-benzyl-N-(4,5-dihydro-imidazol-2-yl)-O-methylhydroxylamine hydrochloride were prep'd. and their structure was det'd. by IR and NMR spectroscopic data as well as X-ray anal. of the imidazolinium azide salt of one of the compds. Binding evaluation for both  $\alpha$ -1- and  $\alpha$ -2-adrenergic receptors in rat brain preps. of these compds. and previously described  $\alpha$ -1,3,5-trihydroxy-2-aryl-imidazolinium salts 11a-d, N-(4,5-dihydroimidazol-2-yl)-1,3,5-trihydroxy-2-aryl-imidazolinium salts 11a-d, N-(4,5-dihydroimidazol-2-yl)-benzimidazoles, and N-(4,5-dihydroimidazol-2-yl)-indoles was performed. Among the compds. tested, 2-[(2-amino-4,5-dichlorophenyl)imino]imidazolinium chloride showed highest binding affinity to  $\alpha$ -1,2-adrenoreceptors ( $K_i = 30$  nM).  
 IT 330685-57-9  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
 (synthesis, structure, and binding of imidazolines to brain  $\alpha$ -1- and  $\alpha$ -2-adrenergic receptors)  
 RN 330685-57-9 CAPLUS  
 CN 1H-imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl


REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L16 ANSWER 5 OF 10 CAPLUS COPYRIGHT 2002 ACS  
 ACCESSION NUMBER: 2000:138409 CAPLUS  
 DOCUMENT NUMBER: 132:260196  
 TITLE: Prediction of distribution coefficients from structure. Comparison of calculated and experimental data for various drugs  
 AUTHOR(S): Tsantili-Kakoulidou, A.; Panderi, I.; Piperaki, S.; Caimzadia, F.; Darvas, F.  
 CORPORATE SOURCE: Department of Pharmacy, University of Athens, Athens, 157 71, Greece  
 SOURCE: European Journal of Drug Metabolism and Pharmacokinetics (1999), 24(3), 205-212  
 CODEN: EUDPD2; ISSN: 0378-7966  
 PUBLISHER: Medecine et Hygiene  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB The efficiency of the program PrologD to predict distribution coeffs. (D) at any pH and pairing ion concn. has been tested using exptl. logD values for various drugs measured under std. conditions of buffers and ionic strength. Clonidine derivs., fluoroquinolones and  $\beta$ -blockers were included as particular pharmacol. classes within the testing data set. Calcns. were performed using the three logP estn. options implemented in the program. PrologD proved to be very efficient and can be of great advantage in drug research. Prediction patterns and correlations between exptl. and calcd. data indicate acceptable results for more than 80% of the data. In addn., comparable studies using the different options permitted suggestions for the more suitable logP estn. method in respect of the particular classes of compds.  
 IT 63346-74-7  
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
 (comparison of calcd. and exptl. data for various drugs in prediction of distribution coeffs. from structure)  
 RN 63346-74-7 CAPLUS  
 CN 1H-imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L16 ANSWER 4 OF 10 CAPLUS COPYRIGHT 2002 ACS  
 ACCESSION NUMBER: 2000:142505 CAPLUS  
 DOCUMENT NUMBER: 132:330832  
 TITLE: Three-dimensional molecular field analyses of octopamine agonists and antagonists for the locust neuronal octopamine receptor class 3  
 AUTHOR(S): Hirashima, A.; Nagata, T.; Pan, C.; Kuwano, E.; Taniguchi, E.; Eto, M.  
 CORPORATE SOURCE: Graduate School, Division of Bioresource and Bioenvironmental Sciences, Kyushu University, Fukuoka, Japan  
 SOURCE: Journal of Molecular Graphics & Modelling (2000), Volume Date 1999, 17(3/4), 198-206  
 CODEN: JMGHFI; ISSN: 1093-3263  
 PUBLISHER: Elsevier Science Inc.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB The quant. structure-activity relationship (QSAR) of a set of 70 octopamine agonists and 20 antagonists against octopamine receptor class 3 (OAR3) in locust nervous tissue was analyzed by mol. field anal. (MFA). MFA of these compds. evaluated effectively the energy between a probe and a mol. model at a series of points defined by a rectangular grid. Contour surfaces for the mol. fields are presented. These results provide useful information in the characterization and differentiation of octopamine receptor types and subtypes.  
 IT 63346-74-7  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
 (three-dimensional mol. field analyses of octopamine agonists and antagonists for locust neuronal octopamine receptor class 3)  
 RN 63346-74-7 CAPLUS  
 CN 1H-imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L16 ANSWER 6 OF 10 CAPLUS COPYRIGHT 2002 ACS  
 ACCESSION NUMBER: 1997:669819 CAPLUS  
 DOCUMENT NUMBER: 127:274156  
 TITLE: Neurotransmitter-receptors as targets for new insecticides  
 AUTHOR(S): Roeder, T.; Degen, J.; Dyczkowski, C.; Gewecke, M.  
 CORPORATE SOURCE: Zoologisches Institut, Universität Hamburg, Neurophysiologie, Hamburg, D-20146, Germany  
 SOURCE: New Strategies in Locust Control (1997), 219-223.  
 Editor(s): Krall, S.; Pevelling, R.; Ba Diallo, D.  
 Birkhauser: Basel, Switz.  
 CODEN: 65EDA4  
 CONFERENCE: Conference  
 LANGUAGE: English  
 AB The locust neuronal octopamine receptor is believed to be an ideal target for highly specific insecticides. The authors characterized a no. of high affinity agonists of this receptor subtype. Using structure-activity relationships, the authors were able to optimize the structure of these compds. in terms of their affinities. A variety of these compds. show a high degree of specificity for insect octopamine receptors vs. vertebrate adrenergic receptors. The high affinity together with the high degree of specificity makes compds. such as the phenyliminoimidazolidines ideal starting points for the development of new insecticides.  
 IT 63346-74-7, NC 20  
 RL: BUU (Biological use, unclassified); PRP (Properties); BIOL (Biological study); USES (Uses)  
 (affinity for locust neuronal octopamine receptor)  
 RN 63346-74-7 CAPLUS  
 CN 1H-imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI) (CA INDEX NAME)




L16 ANSWER 7 OF 10 CAPLUS COPYRIGHT 2002 ACS


ACCESSION NUMBER: 1995:303934 CAPLUS  
 DOCUMENT NUMBER: 122:77274  
 TITLE: Pharmacology of the octopamine receptor from locust central nervous tissue (OAR3)  
 AUTHOR(S): Roeder, Thomas  
 CORPORATE SOURCE: Zool. Inst., Univ. Hamburg, Hamburg, D-20146, Germany  
 SOURCE: British Journal of Pharmacology (1995), 114(1), 210-16  
 CODEN: BJPCRM; ISSN: 0007-1188  
 PUBLISHER: Stockton  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB The present study characterized highly effective agonists from different classes of compds. for the neuronal octopamine receptor (OAR3) of the migratory locust (*Locusta migratoria* L.). Biogenic amines and phenyliminoimidazolidines (PIIs) were employed for the study of structure-activity relationships. The highest affinity PIIs were predominantly those with the substitutions at the positions 2 and 4 of the phenolic ring (e.g. NC 7,  $K_1 = 0.3$  nM, NC8,  $K_1 = 0.81$  nM). Substitutions at these positions always had pos. effects on the affinity of the resp. agonists. Substitutions at the positions 3, 5 and 6, however, always had neg. effects on the affinity. At the position one of the phenolic ring, heterocyclic substituents are preferred. Some PIIs had a more than 30 times higher affinity for OARs than for  $\alpha_1$ -adrenoceptors which are the vertebrate homologues of the insect octopamine receptors. The only non-PII with subnanomolar affinity was the aminooxazoline deriv. AC 6 ( $K_1 = 0.92$  nM). A variety of substances with known insecticidal activity such as chlordimeform, demethylchlordimeform, amitraz or AC 6 had high affinity for the locust neuronal octopamine receptor.

IT 63346-74-7, NC 20  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study) (structure-activity relationship of agonists for locust neuronal octopamine receptor)  
 RN 63346-74-7 CAPLUS  
 CN 1H-Imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI) (CA INDEX NAME)



L16 ANSWER 8 OF 10 CAPLUS COPYRIGHT 2002 ACS (Continued)



L16 ANSWER 8 OF 10 CAPLUS COPYRIGHT 2002 ACS


ACCESSION NUMBER: 1985:593431 CAPLUS  
 DOCUMENT NUMBER: 103:193431  
 TITLE: Phenyliminoimidazolidines. Characterization of a class of potent agonists of octopamine-sensitive adenylyl cyclase and their use in understanding the pharmacology of octopamine receptors  
 AUTHOR(S): Nathanson, James A.  
 CORPORATE SOURCE: Dep. Neurol., Harvard Med. Sch., Boston, MA, 02114, USA  
 SOURCE: Mol. Pharmacol. (1985), 28(3), 254-68  
 CODEN: MOPMAJ; ISSN: 0026-895X  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB Approx. 30 substituted phenyliminoimidazolidines (PII) were examd. for agonist and antagonist effects on the highly enriched and specific octopamine (O)-sensitive adenylyl cyclase (AC) present in the firefly light organ, as well as on ACs present in other invertebrate and vertebrate tissues. Several derivs. were extremely active and some had potencies exceeding those of any previously described agonists of O-sensitive AC. Stimulation by the potent PIIs was reversible, nonadditive to that caused by O, and could be antagonized by antagonists such as cyproheptadine, phentolamine, and propranolol. The inhibitory consts. agreed well with those for inhibiting O stimulation. Certain PII derivs. acted as partial agonists and some as antagonists of O stimulation. Structure-activity relationships revealed, among other things, that short-chain alkyl substitution in the 2- and 6-Phe positions enhanced activity, as did further substitution of 4-halo, 4-Me, or 4-hydroxy substituents. 4-Amino or N-alkyl substitution decreased activity. Structurally related benzyliminoimidazoline derivs. such as tolazoline and naphazoline were partial O agonists, generally less active than the PIIs. Comparison, in 3 invertebrate species, of the effects of the PIIs and 2 other chem. classes of O agonists demonstrated clearcut differences in species responsiveness. Other comparative studies revealed that the agonist activity of the potent PIIs was specific for tissues contg. an O-sensitive AC; ACs activated by dopamine or by  $\beta_1$ - or  $\beta_2$ -adrenergic agonists were unaffected by these compds. The active PIIs affected a class of O receptors distinct from mammalian  $\alpha_1$ - or  $\alpha_2$ -adrenergic receptors. These O receptors also appeared distinct from mammalian 5-HT<sub>1</sub> and 5-HT<sub>2</sub> receptors. Correlative physiol. studies in insects revealed that the active PIIs mimicked O and were potent activators of light emission in the firefly light organ. The PIIs also cause disruption of motor and feeding behavior in tobacco hornworms, leading to insect death, an effect which was markedly potentiated by phosphodiesterase inhibitors.

IT 63346-74-7  
 RL: BIOL (Biological study) (octopamine-sensitive adenylyl cyclase activation by, in light organ of firefly, structure in relation to)  
 RN 63346-74-7 CAPLUS  
 CN 1H-Imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI) (CA INDEX NAME)


L16 ANSWER 9 OF 10 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1977:447931 CAPLUS  
 DOCUMENT NUMBER: 87:47931  
 TITLE: Structure-activity relations and problems related to the mechanism of action of clonidine  
 AUTHOR(S): Rouot, Bruno; Leclerc, Gerard; Wermuth, Camille-Georges; Miesch, Francois; Schwartz, Jean  
 CORPORATE SOURCE: Fac. Pharm., Univ. Louis-Pasteur, Strasbourg, Fr.  
 SOURCE: J. Pharmacol. (1977), 8(1), 95-106  
 CODEN: JNPHAG  
 DOCUMENT TYPE: Journal  
 LANGUAGE: French  
 GI



AB All 26 clonidine analogs (I) studied showed peripheral  $\alpha_1$ -sympathomimetic activity, with IFS 56 (I; R = 2,3-dichloro) [15327-44-3] having the greatest hypertensive effect in demedullated rats. The results correlated with Es (steric const.) and F (sum of the field effect of the substituents) of the Hansch equation (1971). No such correlation was obsd. for hypotensive activity in intact rats. Compds. which had hypotensive activity also had high  $\alpha_1$ -sympathomimetic activity, local anesthetic activity (rabbit cornea), and similar lipophilicity. However, the results were not sufficient to conclude that  $\alpha_1$ -adrenergic mechanisms are involved in the hypotensive effects of clonidine and its analogs.

IT 63346-74-7  
 RL: BAC (Biological activity or effector, except adverse); BIOL (Biological study) (hypotensive and  $\alpha_1$ -sympathomimetic activity of)  
 RN 63346-74-7 CAPLUS  
 CN 1H-Imidazol-2-amine, N-[2,6-bis(1-methylethyl)phenyl]-4,5-dihydro- (9CI) (CA INDEX NAME)



L16 ANSWER 10 OF 10 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1976:456532 CAPLUS

DOCUMENT NUMBER: 85:56532

TITLE: Clonidine and related analogues. Quantitative

correlations

AUTHOR(S): Rouot, Bruno; Leclerc, Gerard; Vermuth, Camille G.;

Miesch, Francois; Schwartz, Jean

CORPORATE SOURCE: Fac. Pharm., Univ. Louis Pasteur, Strasbourg, Fr.

SOURCE: J. Med. Chem. (1976), 19(8), 1049-54


CODEN: JMCNAR

DOCUMENT TYPE: Journal

LANGUAGE: English

GI

L16 ANSWER 10 OF 10 CAPLUS COPYRIGHT 2002 ACS (Continued)




AB A series of 22 derivs. of clonidine-HCl (I) [4205-91-8] were prepd. by the cyclization reaction of ethylenediamine with an S-methylisothiuronium salt deriv. and the main physicochem. parameters (log P, .DELTA.RM, pKa) detd. Quant. correlations between peripheral .alpha.-mimetic action (pithed rats) and physicochem. parameters pointed out the crit. role of the steric effect of ortho substituents. Attempted quant. correlations between physicochem. parameters and central hypotensive activity were unsuccessful. The mechanism of action of I is discussed.

IT 59465-43-9P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(prepn. and blood pressure response to)

RN 59465-43-9 CAPLUS

CN Benzenamine, N-2-imidazolidinylidene-2,6-bis(1-methylethyl)-,  
monohydrochloride (9CI) (CA INDEX NAME)



● HCl

=> FIL STNGUIDE  
 COST IN U.S. DOLLARS  
 FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
49.43	339.92

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)  
 CA SUBSCRIBER PRICE

SINCE FILE	TOTAL
ENTRY	SESSION
-6.20	-7.44

FILE 'STNGUIDE' ENTERED AT 12:32:21 ON 21 AUG 2002  
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT  
 COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE  
 AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.  
 LAST RELOADED: Aug 16, 2002 (20020816/UP).

=> log y  
 COST IN U.S. DOLLARS  
 FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
0.06	339.98

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)  
 CA SUBSCRIBER PRICE

SINCE FILE	TOTAL
ENTRY	SESSION
0.00	-7.44

STN INTERNATIONAL LOGOFF AT 12:32:51 ON 21 AUG 2002